

The Structure of Cones of Matrices

George Phillip Barker

Mathematics Department

University of Missouri—Kansas City

Kansas City, Missouri

and

Raphael Loewy

Mathematics Department

The University of Tennessee

Knoxville, Tennessee

Recommended by Hans Schneider

ABSTRACT

If K is a cone in R_n we let $\Gamma(K)$ denote the cone in the space M_n of $n \times n$ matrices consisting of all A such that $AK \subseteq K$. We show first that $\Gamma(K)$ is indecomposable if and only if K is indecomposable. Next we let $\Gamma(K)^*$ be the dual of $\Gamma(K)$. Then we show that $\Gamma(K) = \Gamma(K)^*$ if and only if K is the image of the nonnegative orthant under an orthogonal transformation.

1. PRELIMINARIES

We shall be dealing with cones and partial orders in the space R_n of n dimensional column vectors and in the space M_n of $n \times n$ real matrices. All topological notions refer to the usual topology in these spaces. The term *cone* is used in a restricted sense, as is seen in the next definition (which applies to both R_n and M_n).

DEFINITION 1.1. Let V be a finite dimensional real vector space. A set $K \subseteq V$ is a cone if

- (1) $x, y \in K, \alpha, \beta \geq 0$ implies $\alpha x + \beta y \in K$;
- (2) K is closed;
- (3) $K \cap (-K) = \{0\}$;
- (4) K^0 , the interior of K , is nonempty.

Frequently, the term cone is used for a set satisfying (1). If it also satisfies (3) it is called *pointed*, and if it satisfies (4) it is called *full*. In finite dimensional spaces a cone is full if and only if it is *reproducing*, that is, $K - K = V$.

A cone K in V induces a partial order if we define $x \leq y$ to mean $y - x \in K$. Then $x < y$ shall mean $y - x \in K$ and $y \neq x$, while $x \ll y$ shall mean $y - x \in K^\circ$. Certain distinguished subsets of K , called *faces*, are central to our work.

DEFINITION 1.2. Let K be a cone in V .

- (1) A subset $F \subseteq K$ is a *face* of K , denoted $F \triangleleft K$, iff
 - (a) $x, y \in F$, $\alpha, \beta \geq 0$ imply $\alpha x + \beta y \in F$,
 - (b) $0 \leq x \leq y$ and $y \in F$ imply $x \in F$.
- (2) If $F \triangleleft K$, then $F - F = V_1$ is a subspace of V called the *span* of F . Its dimension is denoted $\dim F$.
- (3) An *extremal* of K is a face of dimension 1. The set of all extremals of K is denoted $\text{Ext } K$.
- (4) If $S \subseteq K$, then the face generated by S , which is the least face containing S , is denoted by $\varphi(S)$, or $\varphi(x)$ if $S = \{x\}$.

Let K be a cone in R_n . Then the dual cone is the set

$$K^* = \{ y \mid y^T x \geq 0 \text{ for all } x \in K \}.$$

It is easily verified that K^* is a cone in the sense of (1.1).

As is well known, a partial order in R_n generated by a full cone induces in M_n a partial order. In fact, two such partial orders are given in the next definition.

DEFINITION 1.3. Let K be a cone in R_n .

- (1) $\Gamma(K) = \{ A \in M_n \mid AK \subseteq K \}$.
- (2) $\Delta(K) = \text{cl conv} \{ xy^T \mid x \in K, y \in K^* \}$.

Note that in (2) cl denotes the closure and conv the convex hull of the corresponding sets. It is known (cf. Berman and Gaiha [1]) that $\Gamma(K)^* = \Delta(K^*)$.

2. INDECOMPOSABLE CONES

Following Loewy and Schneider [3] we say that a cone K is a *direct sum* of subsets K_1 and K_2 (and we write $K = K_1 \oplus K_2$) if

- (a) $\text{span } K_1 \cap \text{span } K_2 = \{0\}$,
- (b) $K = K_1 + K_2$.

DEFINITION 2.1. The cone K is called *decomposable* iff there exist nonzero subsets K_1 and K_2 such that $K = K_1 \oplus K_2$. Otherwise K is called *indecomposable*.

PROPOSITION 2.1. If $K = K_1 \oplus K_2$ and if $F \triangleleft K$, then there are $F_i \triangleleft K_i$, $F_i \subseteq K_i$, such that $F = F_1 \oplus F_2$.

REMARK. K_i is a face of K , and K_i is a cone in $\text{span } K_i$. Thus we may write $F_i \triangleleft K_i$ (cf. [3]).

Proof. Choose x so that $F = \varphi(x)$. Then x has a unique representation $x = x_1 + x_2$ with $x_i \in K_i$. Let $F_i = \varphi(x_i)$. Then $F_i \triangleleft K_i$ and $\text{span } F_1 \cap \text{span } F_2 = \{0\}$. Now let $y \in F$ and suppose $0 \leq \alpha y \leq x = x_1 + x_2$. We know that $y = y_1 + y_2$ with $y_i \in K_i$ and that $z = (x_1 - \alpha y_1) + (x_2 - \alpha y_2) \in K$. By the uniqueness of representations it follows that $x_i - \alpha y_i \in K_i$, whence $y_i \in F_i$, since $F_i \triangleleft K_i$. Thus $F = F_1 \oplus F_2$. ■

THEOREM 2.1. Let K be a cone in R_n . The following are equivalent:

- (a) K is indecomposable,
- (b) K^* is indecomposable,
- (c) $\Gamma(K)$ is indecomposable,
- (d) $\Delta(K)$ is indecomposable.

LEMMA 2.1. K is indecomposable iff K^* is indecomposable.

Proof. By theorem (3.3) of Loewy and Schneider [3], K is indecomposable iff $I \in \text{Ext } \Gamma(K)$. Also $A \in \Gamma(K)$ iff $A^T \in \Gamma(K^*)$. Hence $I \in \text{Ext } \Gamma(K)$ iff $I \in \text{Ext } \Gamma(K^*)$. ■

LEMMA 2.2. K is indecomposable iff $\Gamma(K)$ is indecomposable.

Proof. Suppose $\Gamma(K)$ is decomposable. Then by Lemma 2.1 $\Gamma(K)^* = \Delta(K^*)$ is decomposable, say $\Delta(K^*) = \Delta_1 \oplus \Delta_2$. Clearly each Δ_i contains at least one rank 1 extremal. For each $y \in \text{Ext } K^*$, $y \neq 0$, put

$$S_i(y) = \{x \mid x \in \text{Ext } K, yx^T \in \Delta_i\}, \quad i = 1, 2.$$

Suppose there is a y such that both $S_1(y) \neq \{0\}$ and $S_2(y) \neq \{0\}$. Then let $F_i = \varphi(S_i) \triangleleft K$. Since each yx^T is in $\text{Ext } \Delta(K^*)$ when $x \in \text{Ext } K$, it follows that $S_1 \cup S_2 = \text{Ext } K$. Thus $F_1 + F_2 = K$. Suppose $z \in \text{span } F_1 \cap \text{span } F_2$. Let $z = z_1 -$

$z_2 = w_1 - w_2$, where $z_1, z_2 \in F_1$, $w_1, w_2 \in F_2$. Then $y(z_1 - z_2)^T = yz_1^T - yz_2^T \in \text{span } \Delta_1$, while $y(w_1 - w_2)^T \in \text{span } \Delta_2$. Thus $yz^T \in \text{span } \Delta_1 \cap \text{span } \Delta_2 = \{0\}$, so $z = 0$ since $y^T \neq 0$. Hence in this case $F = F_1 \oplus F_2$. On the other hand, if there is no such y , then for each $y \in \text{Ext } K^*$ ($y \neq 0$), either $S_1(y) = \{0\}$ or $S_2(y) = \{0\}$. Now let

$$T_1 = \{ y \mid y \in \text{Ext } K^*, y \neq 0, S_2(y) = \{0\} \},$$

$$T_2 = \{ y \mid y \in \text{Ext } K^*, y \neq 0, S_1(y) = \{0\} \}.$$

Then T_1 and T_2 are both nonempty, and we put $G_i = \varphi(T_i)$, $i = 1, 2$. Thus $T_1 \cup T_2 = \text{Ext } K^*$, and so $G_1 + G_2 = K^*$. Suppose $w \in \text{span } G_1 \cap \text{span } G_2$. Then $w = w_1 - w_2 = z_1 - z_2$, where $w_1, w_2 \in G_1$, $z_1, z_2 \in G_2$. Let $x \in \text{Ext } K$, $x \neq 0$. Then

$$(z_1 - z_2)x^T = wx^T = (w_1 - w_2)x^T \in \text{span } \Delta_1 \cap \text{span } \Delta_2.$$

Thus as before $w = 0$ and $K^* = G_1 \oplus G_2$. So by Lemma 2.1, K is also decomposable.

Now suppose $K = K_1 \oplus K_2$. Put

$$\Gamma_{ij} = \{ A \mid A \in \Gamma(K), AK_i \subseteq K_j, AK_l \subset \{0\} \text{ for } l \neq i \}.$$

It is easy to see that $\Gamma(K) = \Gamma_{11} + \Gamma_{12} + \Gamma_{21} + \Gamma_{22}$ and that $\Gamma_{ij} \triangleleft \Gamma(K)$ for each i and j . Put $\Theta = \varphi(\Gamma_{12} + \Gamma_{21} + \Gamma_{22}) \triangleleft \Gamma(K)$. Then $\Gamma_{11} + \Theta = \Gamma(K)$. Let $A \in \Gamma_{11} \cap \Theta$, $x \in K$. Then $x = x_1 + x_2$. Since $A \in \Gamma_{11}$, $Ax_2 = 0$. But also one can find $B_1 \in \Gamma_{12}$, $B_2 \in \Gamma_{21}$, $B_3 \in \Gamma_{22}$ such that $0 \leq A \leq B_1 + B_2 + B_3$. Then $0 \leq Ax_1 \leq B_1x_1 + B_2x_1 + B_3x_1 = B_1x_1 \in K_2$. Thus $Ax_1 \in K_1 \cap K_2 = \{0\}$. Hence $Ax = 0$. Since K is full, $A = 0$. Therefore $\Gamma(K)$ is decomposable. ■

Proof of Theorem 2.1. Since $\Delta(K) = \Gamma(K^*)^*$, the theorem follows from the two lemmas. ■

3. SELF-DUAL CONES

DEFINITION 3.1. Let K be a cone in the real vector space V , and let $\dim V = n$.

- (1) K is *polyhedral* if it has a finite number of extremals.
- (2) K is *simplicial* if it has $n = \dim V$ extremals.
- (3) K is *self-dual* if $K^* = K$.

REMARK 3.1. *It is a well known result that K is simplicial if and only if it induces a lattice order in R_n [2, p. 354].*

DEFINITION 3.2. Subsequently P will denote the nonnegative orthant, that is, the cone of all vectors x such that each component of x is nonnegative.

The next proposition generalizes a lemma communicated to us by B. Levinger at the Auburn Matrix Theory Conference.

PROPOSITION 3.1. *Let I denote the identity matrix. $I \in \Delta(K)$ if and only if K is simplicial.*

Proof. If K is simplicial, then there is a nonsingular $A \in M_n$ such that $P = AK$. Then $\Gamma(K) = A^{-1}\Gamma(P)A$ and $\Delta(K) = A^{-1}\Delta(P)A$. Since $\Gamma(P) = \Delta(P)$, it follows that $\Delta(K) = \Gamma(K)$.

Conversely, suppose $I \in \Delta(K)$. Then there is a sequence $\{\Sigma_n\} \subseteq \Delta(K) \subseteq \Gamma(K)$ of the form

$$\Sigma_n = \sum_{i=1}^{k_n} x_{in} y_{in}^T, \quad x_{in} \in K, \quad y_{in} \in K^*$$

such that $\Sigma_n \rightarrow I$. For any $x, y \in R_n$ we define $\sup\{x, y\}$ as follows. Let

$$\sigma_n = \sum_{i=1}^{k_n} \max\{(y_{in}^T x), (y_{in}^T y)\} x_{in}.$$

We wish to show that $\{\sigma_n\}$ is a bounded sequence. Let μ_1 be a norm which is additive on K [i.e., $x_1, x_2 \in K$ implies $\mu_1(x_1 + x_2) = \mu_1(x_1) + \mu_1(x_2)$]. Let $e \in K^0$, and let μ be the order unit norm corresponding to e with dual norm μ^D and induced matrix norm λ . Then if $y^T \in K^*$ and $A \in \Gamma(K)$, we have

$$\mu^D(y^T) = y^T e, \quad \lambda(A) = \mu(Ae).$$

Since μ_1 and μ are equivalent, there is an $\alpha > 0$ such that for all $x \in R_n$, $\mu_1(x) \leq \alpha \mu(x)$. Since for $y \neq 0$

$$xy^T = [\mu^D(y^T)x][(\mu^D y^T)^{-1}y^T],$$

we may assume that for all i and all n $\mu^D(y_{in}^T) = 1$. Finally we let $M = \max\{\mu(x), \mu(y)\}$. Then we have

$$\begin{aligned}\mu_1(\sigma_n) &\leq \sum_{i=1}^{k_n} |\max\{(\langle y_{in}^T x \rangle, \langle y_{in}^T y \rangle)\}| \mu_1(x_{in}) \\ &\leq M \sum_{i=1}^{k_n} \mu_1(x_{in}) = M \mu_1\left(\sum_{i=1}^{k_n} x_{in}\right) \\ &\leq M \alpha \mu\left(\sum_{i=1}^{k_n} x_{in}\right) = M \alpha \mu\left(\sum_{i=1}^{k_n} (\langle y_{in}^T e \rangle x_{in})\right) \\ &= M \alpha \mu(\Sigma_n e) = M \alpha \lambda(\Sigma_n),\end{aligned}$$

which is bounded, since $\Sigma_n \rightarrow I$ implies $\lambda(\Sigma_n) \rightarrow 1$. Then $\{\sigma_n\}$ is a bounded sequence of vectors, and passing to a subsequence if necessary we obtain a limit $\sigma = \sigma(x, y)$. Note that (taking the limit via the subsequence) we have

$$\begin{aligned}\sigma - x &= \lim_n \sum_{i=1}^{k_n} \max\{(\langle y_{in}^T x \rangle, \langle y_{in}^T y \rangle)\} x_{in} - \lim_n \sum_{i=1}^{k_n} (\langle y_{in}^T x \rangle) x_{in} \\ &= \lim_n \sum_{i=1}^{k_n} [\max\{(\langle y_{in}^T x \rangle, \langle y_{in}^T y \rangle)\} - (\langle y_{in}^T x \rangle)] x_{in} \geq 0.\end{aligned}$$

A similar argument also shows that $\sigma \geq y$.

Now suppose that $z \geq x$ and $z \geq y$. Then $y_{in}^T(z - x) \geq 0$ and $y_{in}^T(z - y) \geq 0$ for all i and n , whence

$$y_{in}^T z - \max\{(\langle y_{in}^T x \rangle, \langle y_{in}^T y \rangle)\} \geq 0.$$

Therefore $z - \sigma \geq 0$.

If σ' is the limit of any other subsequence of the original σ_n , then the above argument remains valid for σ' and its corresponding subsequence. But then both $\sigma \geq \sigma'$ and $\sigma' \geq \sigma$, whence $\sigma = \sigma'$. Thus K is simplicial. ■

THEOREM 3.1. $\Gamma(K) = \Gamma(K)^*$ if and only if $K = QP$ for some orthogonal matrix Q .

We shall obtain the proof as a series of lemmas.

LEMMA 3.1. If $\Gamma(K) = \Gamma(K)^*$, then $K = K^*$.

Proof. If $A \in \Gamma(K)$, $B \in \Gamma(K)^*$, then we may take the inner product as $(B, A) = \text{tr} B^T A$.

Suppose $y \in K^*$ but $y \notin K$. Then there is a $z \in K^*$ such that $z^T y < 0$ and $\forall x \geq 0, z^T x \geq 0$. Since $\Gamma(K) = \Gamma(K)^*$, we have $A \geq 0$ if and only if $A^T \geq 0$. Let $u \in K^0$. Then $zu^T \in \Gamma(K)^* = \Delta(K^*)$ and $uy^T \in \Gamma(K)$, whence $yu^T = (uy^T)^T \in \Gamma(K)$ as well. Thus

$$0 \leq (zu^T, yu^T) = \text{tr} uz^T yu^T = (z^T y)(u^T u) < 0.$$

Therefore $K^* \subseteq K$.

If now $x \in K$ but $x \notin K^*$, pick $w \in K = K^{**}$ such that $w^T x < 0$ and $\forall y \in K^*, y^T w \geq 0$. Let $z \in K^*, z \neq 0$. Then $xz^T \in \Gamma(K)$, whence $zx^T \in \Gamma(K)$ and $zw^T \in \Gamma(K)^*$. Thus

$$0 \leq (zw^T, zx^T) = (z^T z)(w^T x) < 0.$$

Therefore $K \subseteq K^*$ and equality holds. ■

LEMMA 3.2. *If $\Gamma(K) = \Gamma(K)^* = \Delta(K^*)$, then K is simplicial.*

Proof. If $\Gamma(K) = \Gamma(K)^* = \Delta(K^*)$, then $I \in \Delta(K^*)$. Therefore by Proposition (3.1), K^* is simplicial and so is K . ■

PROPOSITION 3.2. *K is a simplicial self-dual cone if and only if $K = QP$ for some orthogonal matrix Q .*

Proof. If $K = QP$, then $K^* = (Q^T)^{-1}P = QP = K$, and so K is self-dual and simplicial. Conversely, suppose $K = K^*$ and $K = AP$, where A is nonsingular. Then $P = P^* = A^T K^* = A^T K$. Thus $A^T A$ is a one to one map of P onto itself. Thus $(A^T A)^{-1} \in \Gamma(P)$ also. Consequently, $A^T A = RD$, where R is a permutation matrix and D is a matrix with positive diagonal elements. In particular $A^T A$ is positive definite symmetric. But since $A^T A = RD$ is positive definite, its principal minors are all positive. Thus RD is a diagonal matrix so $R = I$. Let $B = AD^{-1/2}$. Then B maps P onto K as well, and $B^T B = I$. Thus B is orthogonal. ■

Proof of theorem. If $K = QP$ where Q is orthogonal, then as in the proof of Proposition (3.1) we have that $\Delta(K) = \Gamma(K)$. But also $\Delta(K) = \Delta(K^*)$, so $\Gamma(K) = \Gamma(K)^* = \Delta(K^*)$. Conversely, if $\Gamma(K) = \Gamma(K)^*$, then by Lemmas (3.1) and (3.2), K is a self-dual simplex and we are done by Proposition (3.2). ■

REFERENCES

- 1 A. Berman and P. Caiha, A generalization of irreducible monotonicity, *Linear Algebra Appl.* **5** (1972), 29–38.
- 2 I. M. Glazman and Ju. I. Ljubic, *Finite-Dimensional Linear Analysis*, MIT Press, Cambridge, Mass., 1974.
- 3 R. Loewy and H. Schneider, Indecomposable Cones, *Linear Algebra Appl.*, to be published.

Received 22 November 1974